

Fig. 1. Metatarsus of *Nysius senecionis*. Fig. 2. Metatarsus of *N. graminicola*. Fig. 3. Head, lateral view, of *N. senecionis*. Reproduced, with permission, from Kirby (1986).

ACKNOWLEDGEMENTS

We would like to thank Dr Stephen Judd of the Entomology Department at Liverpool Museum for confirming the identification and for his helpful advice, and Peter Kirby for allowing the reproduction of illustrations from his key.

REFERENCES

- Dolling, W. R. 1985. *Key to the European Lygaeidae: Orsillinae* (issued with Heteroptera Study Group Newsletter No. 6, December 1985).
- Kirby, P. 1986. Key to British Nysius (issued with Heteroptera Study Group Newsletter No. 7, May 1987).

Stichel, W. 1957–59. Illustrierte Bestimmungstabellen der Wanzen. II Europa. (Hemiptera: Heteroptera Europae). Privately published, Berlin, Hermsdorf, 4: 87–95.

NOTE ADDED IN PROOF

Additional British records of *Nysius senecionis* (Schilling) (Hemiptera: Lygaeidae) from Hampshire and Kent.—Since the discovery of this new British bug in East Sussex in September 1992 I have kept a sharp look-out for it in other localities. Until August 1996 all attempts to rediscover the species met with failure, but on 8.viii.1996 I found several specimens of *N. senecionis*, including pairs in-cop, on flowers of ragwort *Senecio jacobaea* (L.) growing in sandy soil at Hyde Common near Fordingbridge, South Hampshire, SU1712. On 15.viii.1996 the species was swept in numbers off ragwort at Shortheath Common, SU7736, and in a sandy field near The Warren, near Oakhanger, SU7735, both in North Hampshire. Finally, on 30.viii.1996, I found a single male on the ground near ragwort growing on sandy ground near Halling, West Kent, TQ7063. All indications are that *N. senecionis* is now firmly established in many parts of southern England.—PETER J. HODGE, 8 Harvard Road, Ringmer, Lewes, East Sussex BN8 5HJ.

Editorial comment.—It appears that *Nysius senecionis* may be spreading and even be moderately common in some parts of south-east England (P. Kirby, pers. com.) and hemipterists are urged to examine recently caught *Nysius* specimens. The editor would welcome short notes in support of this possible expansion of the bug's range. RICHARD A. JONES.